



ArthroGenX Cream

Number of treatments: Approximately 1 month supply

INGREDIENTS: Water, Butylene Glycol, Isododecane, Copaifera Officinalis (Balsam Copaiba) Resin, Dimethicone, Emu Oil, Stearic Acid, Salix Alba (Willow) Bark Extract, Sodium Hyaluronate, Commiphora Mukul Resin Extract, Boswellia Serrata Extract, Dimethyl Sulfone (MSM), Cinnamomum Cassia Bark, Cetyl Myristoleate, Angelica Pubescens Root, Curcuma Longa (Turmeric) Root Extract, Cinnamomum Camphora (Camphor) Wood Oil, Zingiber Officinale (Ginger) Root Oil, Retinyl Palmitate, Glyceryl Stearate, PEG-100 Stearate, Cetearyl Oliviate, Sorbitan Oliviate, Dimethicone PEG-7 Cocoate, Triethanolamine, Tetrasodium EDTA, Phenoxyethanol, Caprylyl Glycol, Sorbic Acid, Acrylates/10-30 Alkyl Acrylate Crosspolymer, Polysilicone 11, Vanillyl Butyl Ether.

DIRECTIONS: Apply a small amount to area of discomfort 1 to 2 times daily or as directed by your healthcare practitioner.

REFERENCES:

1. Ehrenpreis S. Pharmacology of enkephalinase inhibitors: animal and human studies. *Acupunct Electrother Res.* 1985; 10:203-208.
2. Webach, MR, Murray MT. Botanical Influences on Illness: A sourcebook of clinical research. *Third Line Press, Tarzana, CA, 2000.*
3. Rall B et al., Boswellic acids and protease activity (s. auch folgende Abstracts). In: *PM 61 (Abstracts of 43rd Ann Congr): 105.* 1995.
4. Singh GB, Singh S, Bani S. Anti-inflammatory actions of boswellic acids. *Phytomedicine* 3(1):81-5, 1996. methane (curcumin), a non-steroidal anti-inflammatory agent. *J Pharm Pharmac* 25: 447-52, 1973.
6. Diehl HW, May EL. Cetyl myristoleate isolated from Swiss albino mice: an apparent protective agent against adjuvant arthritis in rats. *J Pharm Sci* 1994;83:296-9.
7. Diehl HW. Method for the treatment of osteoarthritis. *U.S. Patent #5,569,676.*
8. Diehl HW. Method of treating rheumatoid arthritis. *U.S. Patent #4,113,881.*
9. Diehl HW. Cetyl myristoleate. *U.S. Patent #4,049,824.*
10. Siemandi H. The effect of cis-9-cetyl myristoleate (CMO) and

ARTHRO GENX CREAM

COMBINES BOTANICALS, NATURAL OILS, AND CONNECTIVE TISSUE REPAIR COMPOUNDS FOR HEALTHY JOINT SUPPORT

- Essential Fatty Acids from Emu Oil for anti-inflammatory benefit
- Extracts of Willow Bark, Myrrh, Boswellia, Turmeric, Angelica, Cinnamon, Ginger, and Camphora for broad-spectrum anti-inflammatory action and pain relief
- Supports reduction of inflammatory leukotrienes, prostaglandins, and cytokines, including TNF alpha, COX 2, and the inflammatory components of the LOX family of mediators
- Cetyl Myristoleate for joint lubrication and T cell immune balance
- MSM for anti-inflammatory action and structural repair support
- Hyaluronic Acid for connective tissue healing support

ARTHROGENX CREAM (TOPICAL) is designed for your patients with musculo-skeletal pain and discomfort. It can be used alone or in conjunction with oral products such as: ArthroGenX capsules, Pain X, EPA/DHA, Flax Oil, etc. This product is formulated with long-term results in mind. It will give an immediate sense of warmth and help ease pain, but unlike most topical products it doesn't stop with just analgesic benefits. ArthroGenX cream has key nutrients that assist in reducing inflammatory mediators, feed cell membranes with essential fatty acids, and support connective tissue repair and proper joint lubrication and flexibility. This is a cream designed to support actual repair—not just for temporary symptomatic benefit.

After an injury initially use RICE (Rest, Ice, Compress, and Elevate). ArthroGenX cream is excellent during this stage of treatment. After the initial swelling dissipates continue to use ArthroGenX cream and consider discontinuing the ice treatment and moving toward gentle heat. Low-level heat will enhance circulation to the injured area and promote a more rapid resolution of the symptoms. High heat is not recommended as it could cause further injury to already damaged cells. Orally, PainX is also excellent to work systemically to resolve inflammation and promote a more rapid response to treatment.

REFERENCES:

11. Srivastava KC and Mustafa T. Ginger (*Zingiber officinale*) in rheumatism and musculoskeletal disorders. *Med Hypothesis* 39:342-8, 1992.
12. Balazs EA, Denlinger JL: Sodium hyaluronate and joint function. *J Equine Vet Sci* 1985; 5: 217-28.
13. Balazs EA: The physical properties of synovial fluid and the special role of hyaluronic acid. In: Helfet A, editor. Disorders of the knee. Philadelphia: Lippincott; 1974: 61-74.
14. Kitoh Y, Katsuramaki T, Tanaka H, et al: Efficacy of SL-1010 (sodium hyaluronate with high molecular weight) on experimental osteoarthritis induced by intra-articularly applied papain in rabbits. *Folia Pharmacol Jpn* 1992; 100:67-76.
15. Hascall VC, Hascall GK: Proteoglycans. In: Hay ED, editor: Cell biology of the extracellular matrix. New York: Plenum Press 1981: p 39-63.
16. Kuettner KE: Osteoarthritis: cartilage integrity and homeostasis. In: Klippel JH, Dieppe PA, editors; Rheumatology, St. Louis: Mosby - Year Book Europe Limited, 1994: p 6.1 - 6.16.
17. Sherrie Schatz and Sheree Lewis, Emu Oil: Reexamining a Natural Remedy with Today's Technology. Schatz Publishing, Nardin, OK, 1996.
18. Ghosh P. Experimental Study to Determine the Anti-Arthritic Activity of New Emu Oil Formulation (EMMP) Royal North Shore Hospital of Sydney, Australia and Dr. Michael Whitehouse at University of Adelaide, Australia. 1993
19. European Scientific Cooperative on Phytotherapy. Salicis cortex. Exeter, UK: ESCOP; 1996-1997:2. *Monographs on the Medicinal Uses of Plant Drugs, Fascicule 4.*
20. Chrubasik S, Eisenberg E, Balan E, et al. Treatment of low back pain exacerbations with willow bark extract: a randomized double-blind study. *Am J Med.* 2000;109:9-14.
21. Meier B, Sticher O, Julkunen-Tiitto R. Pharmaceutical aspects of the use of willows in herbal remedies. *Planta Med.* 1988;54:559-560.
22. Chrubasik S, Eisenberg E, Balan E, et al. Treatment of low back pain exacerbations with willow bark extract: a randomized double-blind study. *Am J Med.* 2000;109:9-14.
23. Schmid B, Ludtke R, Selbmann HK, et al. Efficacy and tolerability of a standardized willow bark extract in patients with osteoarthritis: randomized, placebo-controlled, double blind clinical trial [translated from German]. *Z Rheumatol.* 2000;59:314-320

ARTHRO GENX CREAM

COMPLEMENTARY ORAL PRODUCTS:

- ArthroGenX capsules
- PainX
- BioInflammatory Plus *functional food and capsules*
- EPA/DHA gel caps
- Flax Oil gel caps

ArthroGenX Cream	CONTAINS	BrandX (Popular Products)
NO	cooling effect	YES (Predominant ingredient: menthol)
YES	warming effect	MINIMAL (Camphor)
YES	essential fatty acids	NO
YES	anti-inflammatory botanicals	MINIMAL (Llex extract)
YES	joint lubricants	NO
YES	connective tissue healing support	NO
YES	treatment aimed at long term resolution	NO

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.