



Presents for your consideration:
Diabetone™

- ◆ Promotes healthy blood sugar balance
- ◆ Blood sugar level regulation
- ◆ Nutritional support for metabolism function

Diabetone™ is a complete product that addresses blood sugar function and promotes healthy balance and regulation.

The **BioBalance Diet Plan** rounds out our approach to better blood sugar management. While you the physician, may need to ‘fine tune’ each individuals diet plan and exercise plan, the BioBalance Diet Plan should get your patient started in the right direction. The basic idea of the BioBalance Diet Plan is to eat whole foods with high nutritional value and low glycemic values. Low glycemic foods are those that digest slowly and promote more steady blood sugar levels. This helps to minimize large fluctuations in blood sugar levels. Foods high in sugar or carbohydrates are minimized or avoided. Foods high in fiber, which slows down the absorption of starch and carbohydrates, are promoted. Protein and essential fatty acids are promoted. These foods naturally help to maintain a steady blood sugar level.

Gymnema sylvestre is standardized for gymnemic acid content has been shown to enhance pancreatic beta cell activity and reproduction, thereby improving endogenous insulin output.

Vanadium (vanadyl sulfate) is a salt that has demonstrated insulin-like effects on glucose metabolism. Vanadium appears to enhance insulin receptors at tissue receptor sites. Vanadium improves insulin sensitivity in NIDDM patients.

Chromium has been shown to enhance glucose metabolism and transport. Chromium enhances

Diabetone™	Amounts per serving
Serving size	3 capsules
Number of servings per container	30
Gymnema Sylvestre Extract & Leaf (75% Gymnemic acid)	750 mg.
Chromium (polynicotinate)	500 mcg.
Fenugreek Seed 15:1 extract (trigonella foenum-graecum)	300 mg.
Quercetin	200 mg.
Momordica Charantia	150 mg.
Alpha lipoic acid	150 mg.
Bilberry Extract (25% Anthocyanosides)	85 mg.
Inositol	60 mg.
Vanadium (pentoxide aspartate)	25 mg.
Biotin	16,000 mcg.
Rutin	50 mg.
Suggested Dose: Take 1-2 capsules three times per day or as directed by your health care practitioner.	

insulin responsiveness in skeletal muscle and fat cells. Beneficial effects have also been noted on the serum lipids, cholesterol, triglycerides, HDL cholesterol and LDL cholesterol.

Alpha Lipoic Acid improves insulin resistance in NIDDM patients. Alpha Lipoic Acid favorably influences the glucose transport proteins GLUT1 and GLUT4, thereby enhancing glucose uptake and utilization.

Fiber slows gastric emptying and thus slows glucose absorption through the intestinal mucosa. This benefit creates a more stable blood sugar level and favorably enhances insulin responsiveness.

Biotin is a B vitamin that has been shown to have insulin-like effects and enhances glucose uptake at the insulin receptor site.

Fenugreek Seed and Momordica (Charantin) are natural herbal concentrates have demonstrated improved glucose tolerance and improved glycemic control. Both herbs have been shown to reduce elevated serum glucose levels.

Myoinositol acts as an activator of adenosine triphosphate (ATP) energy mechanisms. It is reported to aid in glycogen utilization, fat metabolism (lipotropic effect), and oxidative phosphorylation. These actions benefit diabetics through enhanced

glucose utilization, fat metabolism and improved energy production through efficient ATP production. **Quercetin and Rutin** are flavonoids that have been shown to act as antioxidants protecting lipids from oxidizing (lipid peroxidation). They also quench the peroxide free radical. Studies have shown them to reduce cardiovascular disease and cancer risk. Mast cell stabilization and reduction of release of inflammatory mediators is also a benefit of quercetin and rutin.

Vaccinium myrtillus with its anthocyanidin pigments has been shown to protect delicate optic structures. Vaccinium extracts can help lower the risk of developing diabetic retinopathy and cataracts.

Recommended Foundational Support:

UltraGenesis™ Multivitamin/Mineral Complex (with or without iron): It is recommended that all your patients be on a therapeutic balanced multivitamin/mineral supplement. With this in mind, we recommend UltraGenesis™. This multivitamin/mineral is rich in antioxidants, coenzyme B vitamins, and macro and micro minerals chelated to the best mineral transporters for ease of absorption.

References:

1. Zhang H, Osada K, Maebashi M, et al. A high biotin diet improves the impaired glucose tolerance of long-term spontaneously hyperglycemic rats with non-insulin-dependent diabetes mellitus. *J Nutr Sci Vitaminol.* 1996;42:517-526.
2. Schechter Y, Li J, Meyerovitch J, et al. Insulin-like actions of vanadate are mediated in an insulin-receptor-independent manner via non-receptor protein tyrosine kinases and protein phosphotyrosine phosphatases. *Mol Cell Biochem.* 1995;153:39-47.
3. Halberstam M, Cohen N, Schimovich P, et al. Oral vanadyl sulfate improves insulin sensitivity in NIDDM but not in obese nondiabetic subjects. *Diabetes.* 1996;45:659-666.
4. Boden G, Chen X, Ruiz J, et al. Effects of vanadyl sulfate on carbohydrate and lipid metabolism in patients with non-insulin-dependent diabetes mellitus. *Metabolism.* 1996;45(9):1130-1135.
5. Linday LA. Trivalent chromium and the diabetes prevention program. *Med Hypotheses.* 1997;49:47-49.
6. Jacob S, Hennksen EJ, Schiemann AI, et al. Enhancement of glucose disposal in patients with type 2 diabetes by alpha-lipoic acid. *Arzneimittelforschung.* 1995;45(8):872-874.
7. Jacob S, Streeper RS, Fogt DL, et al. The antioxidant alpha-lipoic acid enhances insulin-stimulated glucose

- metabolism in insulin-resistant rat skeletal muscle. *Diabetes.* 1996;45:1024-1029.
8. Roebuck JR Jr, Hla KM, Chambless L, Fletcher RH. Effects of chromium supplementation on serum high-density lipoprotein cholesterol levels in men taking beta-blockers. *Ann Intern Med.* 1991;115:917-924.
 9. Anderson R, Cheng N, Bryden NA, et al. Beneficial effects of chromium for people with type 2 diabetes. *Diabetes.* 1996;45(Suppl 2):124A.
 10. Anderson RA, Cheng N, Bryden NA, et al. Elevated intakes of supplemental chromium improve glucose and insulin variables in individuals with type 2 diabetes. *Diabetes.* 1997;46:1786-1791.
 11. Rivellesse A, Giacco A, Genovese S, et al. Effect of dietary fibre on glucose control and serum lipoproteins in diabetic patients. *Lancet.* 1980;2:447-449.
 12. Krezowski PA, Nurrall FQ, Gannon, MC, Bartosh NH. The effect of protein ingestion on the metabolic response to oral glucose in normal individuals. *Am J Clin Nutr.* 1986;44:847-856.
 13. Guevin N, Jacques H, Nadeau A, Galibors I. Postprandial glucose, insulin, and lipid responses to four meals containing unpurified dietary fiber in non-insulin-dependent diabetes mellitus (NIDDM), hypertriglyceridemic subjects. *J Am Coll Nutr.* 1996;15(4):389-396.
 14. Feldman N, Norenberg C, Voet H, Manor E, Berner Y, Madar Z. Enrichment of an Israeli ethnic food with fibres and their effects on the glycemic and insulinaemic responses in subjects with non-insulin-dependent diabetes mellitus. *Br J Nutr.* 1995;74:681-688.
 15. Jacob S, Ruus P, Hermann R, et al. Oral administration of RAC-lipoic acid modulates insulin sensitivity in patients with type-2 diabetes mellitus: a placebo-controlled pilot trial. *Free Rad Biol Med.* 1999;27:309-314.
 16. Estrada DE, Ewart HS, Tsakiridis T, et al. Stimulation of glucose uptake by the natural coenzyme alpha lipoic acid/thiolic acid. *Diabetes.* 1996;45:1798-1804.
 17. Evans Gw, Bowman TD. Chromium picolinate increases membrane fluidity and rate of insulin internalization. *J Inorg Biochem.* 1992;46:243-250.
 18. French RJ, Jones PJH. Role of vanadium in nutrition: metabolism, essentiality and dietary considerations. *Life Sci.* 1993;52:339-346.

These statements have not been evaluated by the Food and Drug Administration. These products are not intended to diagnose, treat, cure, or prevent any disease.

**For Quality and Value without
Compromise**



Is the logical choice!

